

Effect of Recombinant Adenovirus-p53 Combined With Radiotherapy on Long-Term Prognosis of Advanced Nasopharyngeal Carcinoma

JCO February 10, 2009 vol. 27 no. 5 799-804

Jian-ji Pan, Shan-wen Zhang, Chuan-beng Chen, Shao-wen Xiao, Yan Sun, Chang-qin Liu, Xing Su, Dong-ming Li, Gang Xu, Bo Xu, and You-yong Lu

Corresponding author: Shan-wen Zhang, MD, Department of Radiotherapy, Peking University School of Oncology, Beijing Cancer Hospital and Institute, 52 Fucheng Rd, Haidian District, Beijing 100036, People's Republic of China; e-mail: zhangshw4641@sina.com.

ABSTRACT

Purpose

To centrally assess the safety, efficacy, and 6-year follow-up of recombinant adenovirus-p53 (rAd-p53) combined with radiotherapy (RT) for patients with nasopharyngeal carcinoma (NPC).

Patients and Methods

A randomized controlled clinical study on rAd-p53 combined with RT in 42 patients with NPC was compared with a control group of 40 patients with NPC treated with RT alone. In the group receiving rAd-p53 combined with RT, rAd-p53 was intratumorally injected once a week for 8 weeks. Concurrent RT (70 Gy in 35 fractions) was given to the nasopharyngeal tumor and neck lymph node. Patients and tumors were monitored for adverse events and responses.

Results

rAd-p53 – specific p53 mRNA was detected in postinjection of rAd-p53 biopsies from 16 (94.1%) of 17 patients. Upregulation of p21/WAF1 and Bax and downregulation of vascular endothelial growth factor were observed in postinjection tumor biopsy. Complete response rate in the group receiving rAd-p53 combined with RT was observed at 2.73 times that of the group receiving RT alone (66.7% v 24.4%). Six-year follow-up data showed that rAd-p53 significantly increased the 5-year locoregional tumor control rate by 25.3% for patients with NPC treated with irradiation ($P = .002$). The 5-year overall survival rate and 5-year disease-free survival rate of the group receiving rAd-p53 combined with RT were 7.5% ($P = .34$) and 11.7% ($P = .21$) higher than those of the group receiving RT alone. No dose-limiting toxicity or adverse events appeared, except for transient fever after rAd-p53 administration.

Conclusion

In patients with NPC, rAd-p53 was safe and biologically active. Our results indicated that rAd-p53 improves radiotherapeutic tumor control and survival rate in patients with NPC.

INTRODUCTION

Nasopharyngeal carcinoma (NPC) is a cancer unique in the world, and occurs predominantly in southern China. The incidence rate of NPC has been as high as 25 to 50 per 100,000 persons in Guangdong Province, a southern area of China.¹⁻³ Radiotherapy (RT) has been the gold standard of therapy for all stages of NPC for many decades in China, because of anatomic constraints and an intermediate degree of radiosensitivity. A large number of trials in patients with locally advanced NPC have compared concurrent chemotherapy plus RT versus RT alone, but there is uncertainty about the magnitude of survival benefits. Some trials of concomitant chemoradiotherapy failed to detect any improvement on distant failure and survival in Chinese patients with advanced NPC; survival in patients with stages III and IV (M0) disease remained poor.⁴ The 10-year overall survival rate for patients with NPC was 40% to 50%. Patients with stages I and II disease exhibited 10-year overall survival rates of 70% to 80% and 50% to 60%, respectively. Unfortunately, the diagnosis of NPC in most patients occurred at stages III and IV (M0) advanced disease. Current treatments for advanced NPC have been disappointing. The 10-year overall survival rates for patients with stages III and IV (M0) were 30% to 40% and 10% to 20%, respectively. More than 50% of these patients with advanced disease experienced relapse or developed metastasis within 3 years after treatment.¹⁻³

The P53 gene, widely regarded as the genome guardian of cells, plays a key role in cell cycle control, apoptosis, and inhibition of tumor cell proliferation. Wild-type P53 promotes cell cycle arrest and apoptosis of tumor cells after irradiation, but mutated P53 abrogates this response, and induces resistance to radiation.^{5,6} A mutation in the P53 gene occurs in 30% to 70% of patients with NPC, which contributes to recurrence, metastasis, and incurability with regard to irradiation. In patients with NPC, the local recurrence rate is 20% to 50%, and the distant metastasis rate is 20% to 30%.⁷⁻¹⁰ Foundational studies have demonstrated that recombinant adenovirus-p53 (rAd-p53) gene transduction restores P53 function in cancer cells, and increases radiosensitivity of NPC cells in vitro and in vivo. Administration of rAd-p53 could be used in adjuvant therapy with RT.^{11,12} Numerous clinical trials of rAd-p53 gene therapy have initially targeted head and neck squamous cell carcinoma (HNSCC), and rAd-p53 has been demonstrated to be safe, feasible, and effective in HNSCC.^{13,14} Pretreatment experiments showed that expressions of p53 start 6 hours after infection of rAd-p53, increase 24 hours later, reach a pinnacle after 48 to 72 hours, and decrease to disappearance 120 hours later in vitro and in vivo. Therefore, in the clinical trial, 48 to 72 hours after intratumoral injection of rAd-p53 (Genticine; China Shenzhen SiBiono GeneTech Co Ltd, Shenzhen, People's Republic of China), RT was

performed. Injections of rAd-p53 once a week were suitably combined with five fractions of irradiation per week.¹⁵

PATIENTS AND METHODS

P53 Gene Agent and Usage

rAd-p53, a P53 gene agent, is a recombinant replication-incompetent human serotype 5 adenovirus, in which the E1 region is replaced by a human wild-type P53 expression cassette. rAd-p53 limits the infectivity of the virus to only one cycle, and the adenovirus genome does not integrate into host genome DNA, and does not impair any normal cells. rAd-p53 was stored at -20° C in concentrations of 1_10E12 virus particles/mL. rAd-p53 solution was thawed and diluted moderately in physiologic saline according to tumor size within 0.5 hour of use.

Study Design and Patients

This was a prospectively randomized controlled clinical trial. The trial – including protocols and informed consent process – was approved by the Chinese State Food and Drug Administration and Beijing Hospital Ethics Committee on September 12, 2001. The inclusion criteria were as follows. Patients were 18 to 80 years old. They had a histologic diagnosis of nasopharyngeal squamous cell carcinoma, with measurable focus in nasopharynx and neck by CT image and no distant metastasis. Disease was clinically staged according to the fifth edition of the International Union Against Cancer TNM staging system (1997).³ Initial treatment with RT was judged to be the accepted standard of care. Patients had to have a life expectancy of at least 6 months, and a Karnofsky performance score of at least 70. Patients were required to have adequate bone marrow function (WBC count $\geq 4.0 \times 10^9/L$; hemoglobin $\geq 7g/L$; platelet count $\geq 70 \times 10^9/L$) and adequate liver and renal function (AST, ALT, blood urea nitrogen, and creatinine ≤ 1.5 times the upper limit of normal). The exclusion criteria were as follows. Pregnant or nursing women, and patients with uncontrolled serious infections or with serious heart and lung failure, were excluded. From October 2001 to May 2003, a total of 82 patients with NPC were referred to the two participating cancer centers, Beijing Cancer Hospital (Beijing, People's Republic of China) and Fujian Province Cancer Hospital (Fuzhou, People's Republic of China). Patients with NPC who met the inclusion criteria signed the informed consent forms, and were assigned to one of two groups by the randomized table: rAd-p53 combined with RT, or RT alone (control group). Only one tumor site was selected per patient and tautologically injected. The patients receiving rAd-p53 combined with RT received an intratumoral injection of rAd-p53 at 1_10E12 virus particles/mL once a week (on Friday) for 8 weeks per tumor, which was done directly or guided by nasopharyngeal endoscope for nasopharyngeal cavity tumor, or ultrasound for neck node. In the group receiving rAd-p53 combined with RT, there were 22 injection sites in the nasopharynx, and 20 injection sites in the neck. Irradiation was started 2 days after the first injection. For both groups, the same irradiation technique was administered to either the nasopharyngeal tumor or neck lymph node, with the

conventional fractionation of 2 Gy, once a day weekly from Monday to Friday, with a total radiation dose of 70 Gy in 35 fractions by high energy (~ 6 MV) accelerator photons. To assess response, the two largest orthogonal diameters, A and B of nasopharyngeal tumor or neck lymph node, were measured according to computed tomography image at pretreatment and three times post-treatment: 40 Gy dose time point (end of 4th week), 70 Gy dose time point (end of 7th week), and validation time point (2 months after treatment). The tumor size and tumor shrinkage rate were calculated. Immediate tumor response rates, such as complete response (CR), partial response, stable disease, and progressive disease rates, were determined according to the WHO's evaluation standard of solid tumor treatment by tumor shrinkage rate. A comparative study was also performed on the immediate response rate at 40 Gy dose, 70 Gy dose, and validation time points and long-term survival between the two groups.

Reverse Transcriptase Polymerase Chain Reaction and Immunohistochemistry

After intratumoral injection of rAd-p53, the adenoviral particle infects tumor target cells and delivers the adenoviral genome carrying the therapeutic P53 gene to the cell nucleus for transcription. Biopsies at preinjection and 48 hours after the first intratumoral injection of rAd-p53 were assessed for p53 protein and p53 target gene P21, Bax, and vascular endothelial growth factor (VEGF; the downstream p53-transactivated gene) by immunohistochemical staining (IHC) in 50 couple available samples of the 25 patients. IHC staining of paraffin-embedded tissue was performed according to standard methods. Sections in which more than 75% of cells had definitive nuclear reactivity were scored 4+; 50% to 75%, 3+; 25% to 50%, 2+; 5% to 25%, 1+; and fewer than 5%, 0. Biopsies were assessed for exogenous p53 mRNA by reverse transcriptase polymerase chain reaction (RT-PCR) according to previous methods in 17 couple useful comparable samples from the 25 patients.

Adverse Events

Patients were monitored for adverse events. Toxic and adverse events were assessed as light (grade 1), mild (grade 2), serious (grade 3), and life threatening (grade 4) according to the WHO's evaluation standard for adverse events. Particular attention was paid to body temperature.

Statistical Methods

All data were statistically analyzed using SPSS11.5 statistical software (SPSS Inc, Chicago, IL). The primary endpoints of locoregional control, distant metastasis, disease-free survival, and overall survival were estimated using the Kaplan-Meier statistical method. Univariate analysis was used to identify prognostically significant variables for these endpoints. Differences between survival curves were assessed with the log-rank technique; the method was used to evaluate differences between portions. All reported P values resulted from two-sided statistical tests. All P values less than .05 were considered significant.

RESULTS

Detection of exogenous P53 gene expression in tumor biopsy.

rAdp53 – specific p53mRNA was detected by RT-PCR analyses of tissue samples in 16 (94.1%) of 17 assessable samples taken 48 hours after intratumoral injection of rAd-p53. Preinjection biopsies were negative for rAd-p53 – specific p53 mRNA. No rAd-p53 – specific p53 mRNA sequence was found in preinjection biopsies by RT-PCR method. Furthermore, IHC-positive staining scores showed that for p53, preinjection level of 1.44 increased to 2.48 post injection (P= .050); for p21, preinjection level of 0.32 increased to 0.88 postinjection (P= .015); for Bax, preinjection level of 0.92 increased to 1.63 postinjection (P= .088); and for VEGF, preinjection level of 2.80 decreased to 1.50 postinjection (P= .308). IHC analysis revealed increasing rates of p53 (48.0%; 12 of 25), p21/WAF1(44.0%; 11 of 25), and Bax (37.5%; nine of 24), but a decreasing rate of VEGF (21.0%; five of 24) in postinjection tumor biopsy compared with preinjection tumor biopsy.

Table 1. Characteristics of Patients and Tumors

Characteristic	Group Receiving rAdp53 Combined With RT (n = 42)		Group Receiving RT Alone (n = 40)	
	No.	%	No.	%
Sex				
Male	29		31	
Female	13		9	
Age, years				
Median	47.4		50.3	
Standard deviation	12.7		11.3	
Range	19-70		18-71	
Histologic type				
Keratinizing	4		3	
Undifferentiated	38		37	
UICC stage				
I	0		0	
II	11	26.2	12	30.0
III	16	38.1	13	32.5
IV (M0)	15	35.7	15	37.5
UICC stage				
T1	0	00.0	2	5.0
T2	24	57.5	25	62.5
T3	9	21.4	4	10.0
T4	9	21.4	9	22.5
UICC stage				
N0	3	7.1	2	5.0
N1	18	42.9	21	52.5
N2	15	35.7	13	32.5
N3	6	14.3	4	10.0

Abbreviations: rAd-p53, recombinant adenovirus-p53; RT, radiotherapy; UICC, International Union Against Cancer.

Patient Response to Treatment

Between October 1, 2001, and May 1, 2003, 82 patients with NPC were randomly assigned to the group receiving rAd-p53 combined with RT (n_42) or the group receiving RT alone (n_40). No patient withdrew from this study. No significant

differences in sex, age, physical status, histologic type, or clinical stage (including T and N stages) were found between the two groups (Table 1). Baseline average tumor size of the group receiving rAd-p53 combined with RT (882.5; standard deviation [SD], 565.5 mm²) was a little larger than that of the group receiving RT alone (842.7; SD, 599.9mm²), but they were not significantly different (P=.787). In both groups, all patients were in or beyond stage II. Proportion of patients in stages III and IV (M0) was 73.8% (31 of 42) for the group receiving rAd-p53 combined with RT, and 70.0% (28 of 40) for the group receiving RT alone. Average tumor shrinkage rates at 40-Gy dose, 70-Gy dose, and validation time points were 66.9%, 88.4%, and 92.0% for the group receiving rAd-p53 combined with RT, and 44.1%, 64.3%, and 75.1% for the group receiving RT alone, respectively. Response rate of the group receiving rAd-p53 combined with RT was higher than that of the group receiving RT alone at each treatment point (P= .01). At validation time point, the CR rate of the group receiving rAd-p53 combined with RT was 2.73 times that of the group receiving RT alone (66.7% v 24.4%; P=.01; Table 2).

Adverse Events

Total data showed that all the values in blood, urine, and stool examination, and liver and renal function remained within normal range before and after treatment in the group receiving rAd-p53 combined with RT. Lung and heart function were normal before and after treatment. This indicates that rAd-p53 was safe and welltolerated in the patients with NPC. Additionally, rAd-p53 administration did not appear to increase the adverse effects caused by radiation treatment.

All fever events were grade 1 (less than 38° C) and grade 2 (38° C to 40° C), and were transient and self-limited. The incidence of fever among the 42 patients receiving rAd-p53 injections was 81.0% (34 of 42), which included grade 1 fever (43.0%) and grade 2 fever (38.0%). Development of fever was observed as early as approximately 3 hours after injection, lasting about 4 hours and then disappearing spontaneously. Only slight pain and discomfort at local injection sites were felt from the repeat injections. No other adverse effects were found during the subsequent 6 years of follow-up.

6-Year Follow-Up Outcome

The date of last follow-up was December 31, 2007, with a median follow-up duration of 61.0 months (range, 6 to 74 months). No patients were lost to follow-up. Locoregional (nasopharynx and/or neck lymphnode) recurrence had occurred in one patient in the group receiving rAd-p53 combined with RT, and in 10 patients in the group receiving RT alone at the last 6-year follow-up. The median time to recurrence was 73 months (95% CI, 70 to 75 months) in the group receiving rAd-p53 combined with RT, and 59 months (95% CI, 50 to 67 months) in the group receiving RT alone (P=.002). One-, 3-, and 5-year recurrence rates were 0.0%, 2.7%, and 2.7% in the group receiving rAd-p53 combined with RT, and 12.5%, 24.3%, and 28.0% in the group receiving RT alone, respectively (P=.002; Fig 1). Distant metastases were found

in 11 patients in both the group receiving rAd-p53 combined with RT and the group receiving RT alone at the 6-year follow-up. The sites involved were bone (four patients), lung (four patients), and liver (three patients) in the group receiving rAdp53 combined with RT; and bone (four patients), lung (four patients), and liver (three patients) in the group receiving RT alone.

The median time to metastasis was 59 months (95% CI, 51 to 67 months) in the group receiving rAd-p53 combined with RT, and 56 months (95% CI, 48 to 65 months) in the group receiving RT alone (P=.760). One-, 3-, and 5-year metastatic rates were 11.9%, 19.1%, and 26.8% in the group receiving rAd-p53 combined with RT, and 15.1%, 28.8%, and 28.8% in the group receiving RT alone, respectively (P=.760; Fig 2). The median survival time was 59 months (95% CI, 52 to 66 months) in the group receiving rAd-p53 combined with RT, and 54 months (95% CI, 46 to 62 months) in the group receiving RT alone. One-, 3-, and 5-year survival rates were 97.6%, 78.6%, and 66.7% in the group receiving rAd-p53 combined with RT, and 95.0%, 65.0%, and 59.2% in the group receiving RT alone, respectively (P=.340; Fig3). The median disease-free survival time was 58 months (95% CI, 51 to 66 months) in the group receiving rAd-p53 combined with RT, and 49 months (95% CI, 40 to 58 months) in the group receiving RT alone. One-, 3-, and 5-year disease-free survival rates were 90.5%, 81.0%, and 68.7% in the group receiving rAd-p53 combined with RT, and 80.0%, 60.0%, and 57.0% in the group receiving RT alone, respectively (P=.21; Fig 4).

Table 2. Response Rate of Group Receiving rAd-p53 Combined With RT Compared With Group Receiving RT Alone

Group	Treatment Time Point, P < .001														
	40 Gy					70 Gy					Validation				
	FN	CR	PR	SD	PD	FN	CR	PR	SD	PD	FN	CR	PR	SD	PD
rAd-p53 + RT	42	3	31	8	0	42	21	20	1	0	42	28	13	1	0
RT	57	0	25	32	0	60	5	36	19	0	41	10	25	6	0

Abbreviations: rAd-p53, recombinant adenovirus-p53; RT, radiotherapy; FN, focus number of tumor; CR, complete response; PR, partial response; SD, stable disease; PD, progressive disease.




Fig 1. Locoregional recurrence rate of group receiving recombinant adenovirus-p53 (rAd-p53) combined with radiotherapy (GRT) compared with that of group receiving radiotherapy alone (RT).




Fig 2. Distant metastasis rate of group receiving recombinant adenovirus-p53 (rAd-p53) combined with radiotherapy (GRT) compared with that of group receiving radiotherapy alone (RT).




Fig 3. Overall survival rate of group receiving recombinant adenovirus-p53 (rAd-p53) combined with radiotherapy (GRT) compared with that of group receiving radiotherapy alone (RT).




Fig 4. Disease-free survival rate of group receiving recombinant adenovirus-p53 (rAd-p53) combined with radiotherapy (GRT) compared with that of group receiving radiotherapy alone (RT).

DISCUSSION

Our results indicate that successful exogenous gene transfer of p53 can be achieved by intratumoral administration of rAd-p53. The probable clinical mechanism for increased cytotoxicity by rAd-p53 involved, in part, p53-induced cell cycle arrest and apoptosis by upregulation of p21 and Bax, and induced antiangiogenesis by

downregulation of VEGF. There is similarity of rAd-p53 to the other rAd-p53 trials in the world. In one study, after rAd5CMVp53 (Advexin, INGN 201; Introgen Therapeutics, Austin, TX) injection, p53 gene transfer and p53 specific transgene expression were detected using PCR analysis in esophageal squamous cell carcinoma tumor biopsy from all patients, and mRNA levels of p53, p21, and mdm2 increased in all but one patient.¹⁶ In two reports of lung cancer patients, as demonstrated in postinjection biopsies using RT-PCR analysis, vector-specific wild-type p53 mRNA sequences¹⁷ and p21 and Bax expression increased significantly after intratumoral delivery of rAd-p53.¹⁸ After intraperitoneal administration of rAd-p53 (SCH 58500) for recurrent ovarian cancer, successful exogenous gene transfer and expression of p53; upregulation of p21, Bax, and mdm2; and downregulation of survivin were observed in postinjection tumor biopsy.¹⁹ Induction of mRNA and protein expression of p53 and the p53 target gene P21 were demonstrated in samples from patients with bladder cancer treated with intravesical instillation of rAd-p53 (SCH 58500).²⁰

The same foundational study showed that VEGF expression correlated with high vascular grade. Angiogenesis is required for the growth and progression of malignancies. However, wild-type p53 expression was inversely associated with VEGF expression, suggesting that wild-type p53 is involved in the suppression of the VEGF gene. rAd-p53 infection markedly inhibited the expression of an angiogenic factor (VEGF), and increased the expression of a novel antiangiogenic factor (brain-specific angiogenesis inhibitor1), resulting in reduced neovascularization in vivo. Thus, the antiangiogenic effect of adenoviral-mediated wild-type p53 overexpression may contribute to the ability of this viral vector to inhibit tumor growth.²¹⁻²⁴ Our results demonstrated that downregulation of VEGF was observed in tumor biopsy tissue treated by intratumoral injection of rAd-p53. Obvious necrosis was found in tumor biopsy tissues of patients who received intratumoral injections of rAd-p53, which may be correlated with rAd-p53 - induced downregulation of VEGF.

In the last 10 years, adenoviral-mediated P53 gene agent has been used in numerous clinical trials for advanced HNSCC worldwide, either alone or in combination with conventional treatments, such as chemotherapy or RT. rAd-p53 seems to act synergistically with conventional treatments such as chemotherapy and/or RT. In addition, this apparent synergy still exists in patients who were resistant to chemotherapy and RT.^{13, 14} rAd-p53 therapy has been demonstrated to be safe, feasible, and efficient, including local transgenic expression and evidence of local tumor regression for patients with HNSCC^{13,14} and for patients with esophageal cancer,¹⁶ lung cancer,^{17,18} ovarian cancer,¹⁹ bladder cancer,²⁰ and so on.

rAd-p53 is known to trigger a strong immune response in patients. After intratumoral injection of rAd-p53, we observed infiltration of many lymphocytes in biopsies of tumor lesions of patients enrolled onto the trial. In this study, we demonstrated that the CR rate of locoregional tumors at validation time point for the group receiving rAd-p53 combined with RT was 2.73 times that of the group

receiving RT alone ($P = .01$). Furthermore, locoregional tumor control was improved after treatment in the clinical CR associated with rAd-p53; at the 6-year follow-up, the group receiving rAd-p53 combined with RT and the group receiving RT alone had developed 5-year locoregional failure rates of 2.7% and 28.0%. Moreover, rAd-p53 improved radiotherapeutic locoregional tumor control by 25.3%, although the differences in survival were not statistically significant. But from a clinical view, achieving a 7.5% increase in the 5-year survival rate and an 11.7% increase in the 5-year disease-free survival rate for patients with advanced NPC should be clinically significant. The statistical insignificance is a result of the small sample size of the trial. In the future, we plan to conduct a trial with a larger sample size to prove the statistical significance of the product. In conclusion, rAd-p53, as a gene therapy agent, can be combined with RT because of its safe and radiosensitized efficacy in patients with NPC, and represents a potential therapeutic implication for patients with advanced cancer.

REFERENCES

1. Chen WZ, Zhou DL, Luo KS: Long-term observation after radiotherapy for nasopharyngeal carcinoma (NPC). *Int J Radiat Oncol Biol Phys* 16:311-314, 1989
2. Lee AW, Poon YF, Foo W, et al: Retrospective analysis of 5307 patients with nasopharyngeal carcinoma treated during 1976-1985: Overall survival and patterns of failure. *Int J Radiat Oncol Biol Phys* 23:261-270, 1992
3. Au JS, Law CK, Foo W, et al: In-depth evaluation of the AJCC/UICC 1997 staging system of nasopharyngeal carcinoma: Prognostic homogeneity and proposed refinements. *Int J Radiat Oncol Biol Phys* 56:413-426, 2003
4. Chua DT, Sham JS, Au GK, et al: Concomitant chemoradiation for stage III-IV nasopharyngeal carcinoma in Chinese patients: Results of a matched cohort analysis. *Int J Radiat Oncol Biol Phys* 53:334-343, 2002
5. Zhang SW, Xiao SW, Lu YY: Role of p53 tumor suppressor gene in radiosensitivity of human gastric carcinoma cell lines [in Chinese]. *Zhonghua Fangshe Zhongliuxue Zazhi* 2:116-120, 1999
6. Lowe SW, Bodis S, McClatchey A, et al: p53 status and the efficacy of cancer therapy in vivo. *Science* 266:807-810, 1994
7. Soussi T: p53 antibodies in the sera of patients with various types of cancer: A review. *Cancer Res* 60:1777-1788, 2000
8. Gallo O, Chiarelli I, Boddi V, et al: Cumulative prognostic value of p53 mutations and bcl-2 protein expression in head-and-neck cancer treated by radiotherapy. *Int J Cancer* 84:573-579, 1999
9. Lung ML, Hu Y, Cheng Y, et al: p53 inactivating mutations in Chinese nasopharyngeal carcinoma. *Cancer Lett* 133:89-94, 1998
10. Ho KY, Kuo WR, Chai CY, et al: A prospective study of p53 expression and its correlation with clinical response of radiotherapy in nasopharyngeal carcinoma. *Laryngoscope* 111:131-136, 2001
11. Li JH, Lax SA, Kim J, et al: The effects of combining ionizing radiation and

- adenoviral p53 therapy in nasopharyngeal carcinoma. *Int J Radiat Oncol Biol Phys* 43:607-616, 1999
12. Li JH, Chia M, Shi W, et al: Tumor-targeted gene therapy for nasopharyngeal carcinoma. *Cancer Res* 62:171-178, 2002
 13. Clayman GL, el-Naggar AK, Lippman SM, et al: Adenovirus-mediated p53 gene transfer in patients with advanced recurrent head and neck squamous cell carcinoma. *J Clin Oncol* 16:2221-2232, 1998
 14. Clayman GL, Frank DK, Brusio PA, et al: Adenovirus-mediated wild-type p53 gene transfer as a surgical adjuvant in advanced head and neck cancers. *Clin Cancer Res* 5:1715-1722, 1999
 15. Zhang SW, Xiao SW, Lu YY: Effect of adenovirus-mediated p53 gene transfer on apoptosis and radiosensitivity of human gastric carcinoma cell lines. *Chin J Cancer Res* 15:14-18, 2003
 16. Shimada H, Matsubara H, Shiratori T, et al: Phase I/II adenoviral p53 gene therapy for chemoradiation resistant advanced esophageal squamous cell carcinoma. *Cancer Sci* 97:554-561, 2006
 17. Schuler M, Rochlitz C, Horowitz JA, et al: A phase I study of adenovirus-mediated wild-type p53 gene transfer in patients with advanced non-small cell lung cancer. *Hum Gene Ther* 9:2075-2082, 1998
 18. Swisher SG, Roth JA, Komaki R, et al: Induction of p53-regulated genes and tumor regression in lung cancer patients after intratumoral delivery of adenoviral p53 (INGN 201) and radiation therapy. *Clin Cancer Res* 9:93-10E1, 2003
 19. Wen SF, Mahavni V, Quijano E, et al: Assessment of p53 gene transfer and biological activities in a clinical study of adenovirus-p53 gene therapy for recurrent ovarian cancer. *Cancer Gene Ther* 10:224-238, 2003
 20. Kuball J, Wen SF, Leissner J, et al: Successful adenovirus-mediated wild-type p53 gene transfer in patients with bladder cancer by intravesical vector instillation. *J Clin Oncol* 20:957-965, 2002
 21. Fujiwara T, Nishizaki M, Tanaka N: Recombinant adenovirus expressing wild-type p53 is antiangiogenic: Implication for lung cancer gene therapy. *Gan To Kagaku Ryoho* 27:1217-1224, 2000
 22. Zhang L, Yu D, Hu M, et al: Wild-type p53 suppresses angiogenesis in human leiomyosarcoma and synovial sarcoma by transcriptional suppression of vascular endothelial growth factor expression. *Cancer Res* 60:3655-3661, 2000
 23. Nishizaki M, Fujiwara T, Tanida T, et al: Recombinant adenovirus expressing wild-type p53 is antiangiogenic: A proposed mechanism for bystander effect. *Clin Cancer Res* 5:10E15-10E23, 1999
 24. Riccioni T, Cirielli C, Wang X, et al: Adenovirus-mediated wild-type p53 overexpression inhibits endothelial cell differentiation in vitro and angiogenesis in vivo. *Gene Ther* 5:747-754, 1998

Acknowledgment

We thank the patients who participated in this clinical trial, without whose help this study would not have been possible. This study was registered in the WHO

International Clinical Trials Registry Platform, Chinese Clinical Trial Register